



Roman Skibiński · Jacek Golak · Kacper Topolnicki ·
Henryk Witała · Evgeny Epelbaum · Hiroyuki Kamada ·
Hermann Krebs · Ulf-G. Meißner · Andreas Nogga

Modern Chiral Forces Applied to the Nucleon–Deuteron Radiative Capture

Received: 7 December 2016 / Accepted: 19 December 2016
© The Author(s) 2017. This article is published with open access at Springerlink.com

Abstract The chiral nucleon–nucleon interaction with semi-local regularization up to the fifth order of chiral expansion is applied to the nucleon–deuteron radiative capture process. Our theoretical approach is based on the formalism of Faddeev equations and the Siegert theorem is exploited to construct the electromagnetic current operator. The very weak dependence of the differential cross section on values of the regularization parameter is observed. This suggests that the improved chiral two-body interaction is a promising starting point to study electromagnetic processes at low energies.

1 Introduction

The recent progress in constructing chiral nuclear interactions resulted in the new version of the nucleon–nucleon (NN) potential [1,2]. This interaction is an improved version of the force derived in earlier works, starting from [3,4], see [5] for extensive review of this approach. The new model presented in [1,2] includes all terms of the chiral expansion up to $N^4\text{LO}$ and benefits from the improved way in which values of the low energy constants in the long range part of the interaction are established. Namely, in Ref. [1,2] the values of these constants are taken directly from the pion-nucleon scattering without additional fine tuning applied in the previous model. Another difference compared to the older approach is the regularization method used. In the

This article belongs to the special issue “The 23rd European Conference on Few-Body Problems in Physics”.

R. Skibiński (✉) · J. Golak · K. Topolnicki · H. Witała
M. Smoluchowski Institute of Physics, Jagiellonian University, 30348 Kraków, Poland
E-mail: roman.skibinski@uj.edu.pl

E. Epelbaum · H. Krebs
Institut für Theoretische Physik II, Ruhr-Universität Bochum, 44780 Bochum, Germany

H. Kamada
Department of Physics, Faculty of Engineering, Kyushu Institute of Technology, Kitakyushu 804-8550, Japan

Ulf-G. Meißner
Helmholtz-Institut für Strahlen- und Kernphysik, Bethe Center for Theoretical Physics, Universität Bonn, 53115 Bonn, Germany

Ulf-G. Meißner · A. Nogga
Institut für Kernphysik, Institute for Advanced Simulation, Jülich Center for Hadron Physics, Forschungszentrum Jülich, 52425 Jülich, Germany

Ulf-G. Meißner · A. Nogga
JARA - High Performance Computing, Forschungszentrum Jülich, 52425 Jülich, Germany

previous model a nonlocal regularization in momentum space has been applied. This means, that the matrix elements of the potential $V: \langle \mathbf{p} | V | \mathbf{p}' \rangle$ have been multiplied by an exponential factor $\exp(-(p^6 + p'^6)/\Lambda^6)$ where \mathbf{p}' and \mathbf{p} are the relative momenta of nucleons in the initial and final state, respectively, $p' = |\mathbf{p}'|$, $p = |\mathbf{p}|$ and $\Lambda \approx 550$ MeV is the regularization parameter. Such a regularization, implemented in the same way for all partial waves, leads to unwanted artifacts in the long-range part of the NN force and does not completely eliminate the unwanted short-range components of the two-pion exchange. Further, the corresponding non-local regularization has been used for the three-nucleon (3N) force [6], what influenced the description of observables in the 3N sector, see [7], and [8] for applications in the deuteron breakup process and [9–11] for the electromagnetic processes including proton–deuteron radiative capture as well as ^2H and ^3He photodisintegration reactions. These works have revealed that the cut-off dependence of the studied there model of nuclear forces is too strong (especially at N^3LO) and precludes precise conclusions about investigated processes.

In the recent model of Refs. [1,2] another regularization method is used. Namely, the long-range part of the interaction in coordinate space is multiplied with the function $f(r) = (1 - \exp(-r^2/R^2))^6$, while contact interactions are regularized using a non-local Gaussian regulator in momentum space. The values of the cutoff parameter R are chosen in the range of 0.8–1.2 fm, however they do not describe the two-nucleon phase shifts equally well - the best description (up to $E_{\text{lab}} = 300$ MeV) is obtained for R of 0.9 and 1.0 fm.

The precise description of 3N systems is more challenging. Even restricting theoretical calculations to the two-body phenomenological forces only, due to the important role of the off-the-energy-shell parts of the interaction, one observes clear model dependence. It was illustrated many times, for example in [12,13], clearly pointing to the necessity of improvement of phenomenological forces employed in these calculations. The chiral interaction has more solid theoretical foundations and in principle should approach the exact theory. However, regulator functions, unavoidable in practical calculations, introduce unwanted uncertainties. While in nucleon–deuteron (Nd) elastic scattering this uncertainty is on the tolerable level for both regularization schemes [6,14], in the Nd radiative capture as well as in the ^2H and ^3He photodisintegration processes the cut-off dependence of predictions based on the nonlocal regularization is huge, even at low energies [9–11].

In this paper we test the regulator dependence of predictions based on the local regularization for the radiative Nd capture process. We briefly summarize our method in Sec. 2 and present results, at two reaction energies, for the differential cross section in Sec. 3. Other observables and more detailed discussion comprising also photodisintegration and weak muon-capture processes can be found in [15].

2 Formalism

Since our approach has been described in detail for example in [16,17], here we only remind the reader of our key equations for the two-body photodisintegration nuclear matrix element N_{Nd}^μ . If the 3N force is neglected, we simply get:

$$N_{Nd}^\mu = \langle \phi_{Nd} | (1 + P) j_{3N}^\mu | \Psi_{\text{bound}}^{3N} \rangle + \langle \phi_{Nd} | P | U^\mu \rangle, \quad (1)$$

$$| U^\mu \rangle = t G_0 (1 + P) j_{3N}^\mu | \Psi_{\text{bound}}^{3N} \rangle + t G_0 P | U^\mu \rangle, \quad (2)$$

where j_{3N}^μ , P , t and G_0 are the electromagnetic current operator, the permutation operator, the t-matrix operator (built from the two-body interaction) and the free 3N propagator, respectively. Further, $|\Psi_{\text{bound}}^{3N}\rangle$ and $|\phi_{Nd}\rangle$ denotes the initial 3N bound state and the final Nd state, respectively. The nuclear matrix element for the radiative Nd capture can be obtained from N_{Nd}^μ by time-reversal symmetry.

The electromagnetic current operator consistent with the strong interaction with the semi-local regularization has not been derived yet (see [18,19] for the first steps in this direction). Thus we follow [16,17] and use the standard single nucleon current operator together with the Siegert theorem to include implicitly many-body contributions to the electromagnetic current.

3 Results and Summary

In Fig. 1 we exemplify the unsatisfactory description of the Nd radiative capture cross section delivered by the nonlocally regularized chiral forces [4]. The cyan (red) band comprises N^2LO (N^3LO) predictions obtained with various values of the cut-off parameters, as suggested in [4]. It is clear that the width of the bands at both orders of the chiral expansion and for both processes presented here (the neutron-deuteron capture at




Fig. 1 The differential cross section $d^2\sigma/d\Omega$ for the $n + d \rightarrow {}^3\text{H} + \gamma$ reaction at $E_n = 9.0\text{ MeV}$ (left) and for the $p + d \rightarrow {}^3\text{He} + \gamma$ reaction at $E_d = 95\text{ MeV}$ (right). Theoretical predictions have been obtained using the chiral forces with nonlocal regularization [4]. The cyan and red bands cover predictions at $N^2\text{LO}$ and $N^3\text{LO}$ obtained with different values of the regularization parameters, respectively. The thick violet dotted curves show predictions based on the AV18 potential. The circles represent data from Ref. [20] (left) and from Ref. [21] (right)




Fig. 2 The differential cross section $d^2\sigma/d\Omega$ for the $n + d \rightarrow {}^3\text{H} + \gamma$ reaction at $E_n = 9.0\text{ MeV}$. The left column shows the convergence of predictions at $R = 0.9\text{ fm}$ with respect to the order of chiral expansion. The double-dotted-dashed green, dashed blue, dotted black, solid red and thick dashed black curves represent predictions at LO, NLO, $N^2\text{LO}$, $N^3\text{LO}$ and $N^4\text{LO}$, respectively. The right panel shows the dependence of the predictions at $N^4\text{LO}$ on the value of the R parameter. The dotted green, thick dashed black, solid black and double-dotted-dashed blue curves correspond to predictions with $R = 0.8, 0.9, 1.0, 1.1$ and 1.2 fm , respectively. Note that the thick dashed black curve shown in both panels represents the same predictions. Also the thick violet dotted curve which represents the AV18 predictions appears in both panels and in the left panel of Fig. 1. The data are from Ref. [20]




Fig. 3 The differential cross section $d^2\sigma/d\Omega$ for the $p + d \rightarrow {}^3\text{He} + \gamma$ reaction at $E_d = 95\text{ MeV}$. Curves as in Fig. 2; the AV18 based predictions are the same as in the right panel of Fig. 1. The data are from Ref. [21]

$E_n = 9\text{ MeV}$ and the proton–deuteron capture at $E_d = 95\text{ MeV}$) is too big to allow one to investigate details of the underlying Hamiltonian.

A big improvement can be observed when the chiral force with the semi-local regularization is used. In Figs. 2 and 3 we show predictions for the same cross sections as in Fig. 1. Now, instead of bands showing the dependence on the value of the regularization parameter R , we present in the right panels separate lines for $R = 0.8, 0.9, 1.0, 1.1$ and 1.2 fm . Since the values $R = 0.9\text{ fm}$ and $R = 1.0\text{ fm}$ give the best description of

the NN system, combining all curves into one band could be somewhat misleading. We observe that all values of R lead to nearly the same values of the cross sections. Moreover, at a lower energy results with extreme values $R = 0.8$ fm (green curve) and $R = 1.2$ fm (blue curve) lie slightly above the remaining, practically overlapping lines. Also at $E_d = 95$ MeV predictions based on $R = 0.8$ fm and $R = 1.2$ fm slightly deviate from the other ones, although the spread of predictions remains small.

In the left panels of Figs. 2 and 3 we show the convergence of predictions with respect to the order of chiral expansion. At both energies predictions obtained at N^3 LO and N^4 LO are practically indistinguishable. They differ from predictions at NLO and N^2 LO, which are close to one another, which is expected due to the appearance of new contact interactions at N^3 LO.

Results presented here and in Ref. [15] obtained with NN forces only confirm a good quality of the chiral force model proposed in Refs. [1,2]. However, the final judgment on quality and usefulness of semi-locally regularized interaction in analysis of electromagnetic three-nucleon reactions could be formulated only after inclusion of three-nucleon forces and consistent electromagnetic current to theoretical analysis. Such work is in progress.

Acknowledgements This work is a part of the LENPIC project. It was financed from the resources of the National Science Center (Poland) under grants DEC-2013/10/M/ST2/00420 and DEC-2013/11/N/ST2/03733. The numerical calculations were done on the computers of the JSC Jülich.

Open Access This article is distributed under the terms of the Creative Commons Attribution 4.0 International License (<http://creativecommons.org/licenses/by/4.0/>), which permits unrestricted use, distribution, and reproduction in any medium, provided you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons license, and indicate if changes were made.

References

1. E. Epelbaum et al., Improved chiral nucleon–nucleon potential up to next-to-next-to-next-to-leading order. *Eur. Phys. J. A* **51**, 5,53 (2015)
2. E. Epelbaum et al., Precision nucleon–nucleon potential at fifth order in the chiral expansion. *Phys. Rev. Lett.* **115**, 12,122301 (2015)
3. E. Epelbaum et al., Nuclear forces from chiral Lagrangians using the method of unitary transformation. 1. Formalism. *Nucl. Phys. A* **637**, 107 (1998)
4. E. Epelbaum et al., The two-nucleon system at next-to-next-to-next-to-leading order. *Nucl. Phys. A* **747**, 362 (2005)
5. E. Epelbaum et al., Modern theory of nuclear forces. *Rev. Mod. Phys.* **81**, 1773 (2009)
6. E. Epelbaum et al., Three nucleon forces from chiral effective field theory. *Phys. Rev. C* **66**, 064001 (2002)
7. H. Witała et al., Calculations of three-nucleon reactions with N^3 LO chiral forces: achievements and challenges. *J. Phys. G: Nucl. Part. Phys.* **41**, 094011 (2014)
8. J. Golak et al., Low-energy neutron–deuteron reactions with N^3 LO chiral forces. *Eur. Phys. J. A* **50**, 177 (2014)
9. R. Skibiński et al., Nucleon–deuteron capture with chiral potentials. *Acta Phys. Polon. B* **37**, 2905 (2006)
10. R. Skibiński et al., The chiral long-range two-pion exchange electromagnetic currents in radiative nucleon–deuteron capture. *Acta Phys. Polon. B* **46**, 159 (2015)
11. D. Rozpędzik et al., Signatures of the chiral two-pion exchange electromagnetic currents in the ^2H and ^3He photodisintegration reactions. *Phys. Rev. C* **83**, 064004 (2011)
12. H. Witała et al., Nd elastic scattering as a tool to probe properties of 3N forces. *Phys. Rev. C* **63**, 024007 (2001)
13. W. Glöckle et al., The three-nucleon continuum: achievements, challenges and applications. *Phys. Rep.* **274**, 107 (1996)
14. S. Binder et al., Few-nucleon systems with state-of-the-art chiral nucleon–nucleon forces. *Phys. Rev. C* **93**, 044002 (2016)
15. R. Skibiński et al., Testing semi-local chiral two-nucleon interaction in selected electroweak processes. *Phys. Rev. C* **93**, 064002 (2016)
16. J. Golak et al., Faddeev calculations of proton–deuteron radiative capture with exchange currents. *Phys. Rev. C* **62**, 054005 (2000)
17. J. Golak et al., Electron and photon scattering on three-nucleon bound states. *Phys. Rep.* **415**, 89 (2005)
18. S. Kölling et al., Two-pion exchange electromagnetic current in chiral effective field theory using the method of unitary transformation. *Phys. Rev. C* **80**, 045502 (2009)
19. S. Kölling et al., Two-nucleon electromagnetic current in chiral effective field theory: one-pion exchange and short-range contributions. *Phys. Rev. C* **84**, 054008 (2011)
20. G. Mitev et al., Radiative neutron capture by deuterium. *Phys. Rev. C* **34**, 389 (1986)
21. W.K. Pitts et al., $^1\text{H}(\text{d}, \gamma)^3\text{He}$ reaction at $E_d = 95$ MeV. *Phys. Rev. C* **37**, 1 (1988)